

Table 1. Preoperative patient demographics and perioperative outcomes

|  | Group I (SCD, n=73)   | Group II (ECD, n=27)  | P value |
|--|-----------------------|-----------------------|---------|
| <b>I. Donor parameter</b>                  |                       |                       |         |
| Age (years)                                | 43 (30.5 – 46)        | 51 (42 – 55)          | < 0.001 |
| Sex (M/F)                                  | 44 / 29               | 25 / 2                | 0.002   |
| Smoking history (pack-years)               | 0 (0 – 6.25)          | 20 (0 – 30)           | < 0.001 |
| Mechanical ventilation (days)              | 5.5 (2.5 – 7.1)       | 5.0 (2.4 – 7.4)       | 0.787   |
| PaO <sub>2</sub> / FiO <sub>2</sub> (mmHg) | 474.1 (417.5 – 521.4) | 437.5 (368.4 – 466.0) | 0.004   |
| <b>II. Recipient parameter</b>             |                       |                       |         |
| Age (years)                                | 49 (37 – 57)          | 55 (43 – 59)          | 0.327   |
| Sex (M/F)                                  | 42/31                 | 14/13                 | 0.611   |
| ABO type (A / B / O / AB)                  | 25 / 21 / 17 / 10     | 10 / 6 / 10 / 1       | 0.306   |
| Disease entity                             |                       |                       | 0.578   |
| Pulmonary fibrosis                         | 33 (45.2%)            | 15 (55.6%)            |         |
| Bronchiectasis                             | 7 (9.6%)              | 0                     |         |
| LAM  | 6 (8.2%)              | 2 (7.4%)              |         |
| BOS  | 9 (12.3%)             | 2 (7.4%)              |         |
| Others                                     | 18 (24.7%)            | 8 (29.6%)             |         |
| Mechanical ventilation                     | 21 (28.8%)            | 9 (33.3%)             | 0.806   |
| ECMO support                               | 11 (15.1%)            | 7 (25.9%)             | 0.245   |
| <b>III. Operative data</b>                 |                       |                       |         |
| Type of transplantation                    |                       |                       | > 0.999 |
| Single                                     | 7 (9.6%)              | 2 (7.4%)              |         |
| Double                                     | 66 (90.4%)            | 25 (92.6%)            |         |
| Cardiopulmonary support                    |                       |                       | 0.063   |
| CPB  | 23 (31.5%)            | 3 (11.1%)             |         |
| ECMO                                       | 41 (56.2%)            | 22 (81.5%)            |         |
| None                                       | 9 (12.3%)             | 2 (7.4%)              |         |
| Immediate postoperative ECMO               | 21 (28.8%)            | 16 (59.3%)            | 0.006   |
| Operation time (mins)                      | 408.0 (363.5 – 464.0) | 407.0 (336.0 – 462.0) | 0.295   |
| <b>IV. Postoperative outcome</b>           |                       |                       |         |
| Re-implantation of ECMO                    | 8 (11.0%)             | 6 (22.2%)             | 0.194   |
| Mechanical ventilation (days)              | 7 (3 – 12)            | 11 (5 – 20)           | 0.445   |
| ICU stay (days)                            | 9 (5.5 – 14.5)        | 12 (8 – 19)           | 0.290   |
| Hospital stay (days)                       | 29 (25.5 – 59.0)      | 31 (24.0 – 45.0)      | 0.051   |
| 30 day mortality                           | 10 (14.1%)            | 2 (7.7%)              | 0.505   |
| Rejection (acute /chronic)                 | 4 / 1                 | 1 / 0                 | > 0.999 |